

Searches for CP , T and CPT violation in $B^0\bar{B}^0$ mixing at $BABAR$

R. Covarelli*

Università degli Studi di Perugia, I-06123 Perugia, Italy

We describe measurements of CP and CPT violation parameters in $B^0\bar{B}^0$ oscillations, performed at $BABAR$ by using $B\bar{B}$ events collected at the $\Upsilon(4S)$ mass peak. These are obtained from two different B -reconstruction methods, namely: searching for two inclusive semileptonic B^0 decays in an event or partially reconstructing one of the B^0 mesons in the $D^{*-}\ell^+\nu_\ell$ channel. The results are in agreement with one another and with the most recent theoretical calculations in the Standard Model scenario.

I. THEORETICAL OVERVIEW

CP violation due to the neutral- B meson oscillation processes can be parameterized in two different scenarios: either in the Standard Model (SM) field theory, with some extensions allowed within the CPT -conservation hypothesis, or, more generally, also taking into account the possibility of CPT violation.

In the SM case, $B^0\bar{B}^0$ mixing is accounted for using a time-dependent perturbation theory that leads to an effective hamiltonian $\mathbf{H}_{eff} = \mathbf{M} - \frac{i}{2}\mathbf{\Gamma}$, that is the 2×2 projection of the actual interaction operator on the B^0 and $\bar{B}^0$ states only. In solving the eigenvalue problem, linear relations are found connecting the B flavor eigenstates to the physical states:

$$\begin{aligned} |B_L\rangle &= p|B^0\rangle + q|\bar{B}^0\rangle, \\ |B_H\rangle &= p|B^0\rangle - q|\bar{B}^0\rangle. \end{aligned} \quad (1)$$

In the Standard Model, the magnitude of the ratio q/p is very nearly unity:

$$\left|\frac{q}{p}\right|^2 \approx 1 - \text{Im}\frac{\Gamma_{12}}{M_{12}}, \quad (2)$$

not just because $|\Gamma_{12}|$ is small, but because the CP -violating quantity $\text{Im}(\Gamma_{12}/M_{12})$ is suppressed by an additional factor $(m_c^2 - m_u^2)/m_b^2 \approx 0.1$ relative to $|\Gamma_{12}/M_{12}|$. When the remaining factors are included, the expectation is $|\text{Im}(\Gamma_{12}/M_{12})| < 10^{-3}$ [1, 2].

The parameter $|q/p|$ is sensitive to New Physics (NP) within a CPT -conserving theory [3]. Assuming 3-generation unitarity of the CKM matrix [4] and tree-level processes dominated by the SM, the CP asymmetry can be modified as follows:

$$\begin{aligned} A_{CP/T} &\simeq 2(1 - |q/p|) = \\ &-\text{Re}\left(\frac{\Gamma_{12}}{M_{12}}\right)^{\text{SM}} \frac{\sin 2\phi_d}{C_d^2} + \text{Im}\left(\frac{\Gamma_{12}}{M_{12}}\right)^{\text{SM}} \frac{\cos 2\phi_d}{C_d^2}, \end{aligned} \quad (3)$$

C_d and ϕ_d being a generic NP amplitude and phase. Within the NP parameter space, $|q/p| - 1$ could be enhanced up to an order of magnitude.

If CPT violation is also allowed, its effects are usually parameterized in terms of an additional complex parameter z in the eigenstate definitions (1):

$$\begin{aligned} |B_L\rangle &= p\sqrt{1-z}|B^0\rangle + q\sqrt{1+z}|\bar{B}^0\rangle, \\ |B_H\rangle &= p\sqrt{1+z}|B^0\rangle - q\sqrt{1-z}|\bar{B}^0\rangle. \end{aligned} \quad (4)$$

Kostelecký [5] first pointed out that such parameterization using a constant value of z is not completely satisfactory, since CPT violation in a field theory implies that Lorentz covariance, too, is violated. A more appropriate parameterization is achieved by setting $z \propto \beta^\mu \Delta a_\mu$, where $\beta^\mu \equiv \gamma_B(1, \vec{\beta}_B)$ is the decaying B -meson four-velocity and Δa_μ is a four-vector containing NP CPT -violating coefficients. We approximate the 4-velocity of each B meson by the $\Upsilon(4S)$ 4-velocity so that z is common to each B in an event; in this way the 3-velocity direction becomes a function of the Earth position with respect to the Universe at the time of the $\Upsilon(4S)$ decay, i.e. of the event's sidereal time. z is then expressed as:

$$z \equiv z(\hat{t}) = z_0 + z_1 \cos(\Omega \hat{t} + \phi) \quad (5)$$

where $\Omega = 2\pi$ rad/sidereal day is the Earth rotation frequency (1 sidereal day ~ 0.997 solar days), $\hat{t}$ is the sidereal time, z_0 and z_1 contain NP coefficients.

All the analyses presented in the following exploit the Δt dependence of the B -pair decay rates in order to extract CP and CPT violation parameters, Δt being the difference between the decay times of the two mesons. The general form is:

$$\begin{aligned} \frac{dN}{d(\Delta t)} &\propto e^{-\Gamma|\Delta t|} \left[\frac{1}{2}c_+ \cosh\left(\frac{\Delta\Gamma\Delta t}{2}\right) + \frac{1}{2}c_- \cos(\Delta m t) \right. \\ &\quad \left. - \text{Res} \sinh\left(\frac{\Delta\Gamma\Delta t}{2}\right) + \text{Im}s \sin(\Delta m t) \right]. \end{aligned} \quad (6)$$

In this equation Δm ($\Delta\Gamma$) is the mass (lifetime) difference between the B^0 physical states, while c_+ , c_- and s depend on the physical parameters $|q/p|$ and z , on the amplitudes of the particular processes under consideration, and the flavor of the B mesons at the time of decay.

II. THE INCLUSIVE DILEPTON METHOD

Inclusive dilepton events, where both B mesons decay semileptonically ($b \rightarrow X\ell\nu$, with $\ell = e$ or μ), represent

*Electronic address: covarelli@slac.stanford.edu

4% of all $\Upsilon(4S) \rightarrow B\bar{B}$ decays and provide a very large sample with which to study T , CP and CPT violation. In the direct semileptonic neutral B decay, the flavor B^0 ($\bar{B}^0$) is tagged by the charge of the lepton ℓ^+ (ℓ^-). This study [6] is performed using a BABAR [7] data sample equivalent to an integrated luminosity of 211 fb^{-1} recorded at the $\Upsilon(4S)$ resonance (232 million $B\bar{B}$ pairs).

Electrons and muons are selected using information from the tracking systems, the electromagnetic calorimeter and the muon chambers. The track momentum is required to be between 0.8 and 2.3 GeV/c and certain dilepton mass ranges are vetoed in order to reject photon conversions, J/ψ and $\psi(2S)$ decays. The separation between *direct* leptons ($b \rightarrow \ell$) and background from the $b \rightarrow c \rightarrow \ell$ decay chain (*cascade* leptons) is achieved with a neural network that combines five discriminating variables: the momenta and opening angle of the two lepton candidates, and the total visible energy and missing momentum of the event.

Since $|q/p|$ is expected to be small, we have determined the possible charge asymmetries induced by charge-dependent differences in the reconstruction and identification of electrons. The charge asymmetry of track reconstruction is measured in the data by comparing tracks reconstructed using only the silicon tracker with those passing the dilepton track selection. The lepton identification efficiencies are measured with a control sample of radiative Bhabha events for electrons. Based on tracking and identification, we fix the charge asymmetry for the direct electrons to $a_e^{dir} = 1.2 \times 10^{-3}$. In the likelihood fit, we float the charge asymmetries a_μ^{dir} and a_μ^{casc} for direct and cascade muons.

We model the contributions to our sample from $B\bar{B}$ decays using different categories of events each represented by a probability density function (PDF) in Δt . The five categories are the following. First, the pure signal events with two direct leptons are 81% of the $B\bar{B}$ events. Then, we consider two categories of cascade decays: those in which the direct lepton and the cascade lepton come from different B decays, and those in which the direct lepton and the cascade lepton stem from the same B decay. In addition, a small fraction of the dilepton events originate from the decay chain $b \rightarrow \tau^- \rightarrow \ell^-$ which tags the B flavor correctly. Finally, the remaining events consist mainly of leptons from the decay of a charmonium resonance. The last component of the dilepton sample originates from non- $B\bar{B}$ events, and has been estimated using off-resonance data. Figure 1 shows the result of the maximum likelihood fit in terms of the projected asymmetry $A_{CP/T}$ between (ℓ^+, ℓ^+) and (ℓ^-, ℓ^-) dileptons and A_{CPT} between (ℓ^+, ℓ^-) dileptons with $\Delta t > 0$ and $\Delta t < 0$.

For $|q/p|$, the most important systematic uncertainties are due to the correction of electron charge asymmetries, while the dominant systematic uncertainties on z are the imperfect knowledge of the absolute scale of the detector and the residual uncertainties in the tracker local alignment.




FIG. 1: (a) $A_{CP/T}$ asymmetry between (ℓ^+, ℓ^+) and (ℓ^-, ℓ^-) . (b) A_{CPT} asymmetry between (ℓ^+, ℓ^-) dileptons with $\Delta t > 0$ and $\Delta t < 0$.

The results for CPT and CP violation parameters are

$$\begin{aligned} |q/p| - 1 &= (-0.8 \pm 2.7_{\text{(stat.)}} \pm 1.9_{\text{(syst.)}}) \times 10^{-3}, \\ \text{Im} z &= (-13.9 \pm 7.3_{\text{(stat.)}} \pm 3.2_{\text{(syst.)}}) \times 10^{-3}, \\ \Delta\Gamma \cdot \text{Re} z &= (-7.1 \pm 3.9_{\text{(stat.)}} \pm 2.0_{\text{(syst.)}}) \times 10^{-3} \text{ ps}^{-1}. \end{aligned}$$

where $\Delta\Gamma \cdot \text{Re} z$ has been determined with the approximation $\text{Re} z \sinh(\Delta\Gamma\Delta t/2) \sim \text{Re} z \cdot \Delta\Gamma \cdot \Delta t/2$ and z is considered as a constant parameter.

A similar Δt fit is then performed, taking the sidereal-time dependence into account. We use the latitude and longitude of the BABAR detector, together with its orientation to determine the Lorentz boost direction of the $\Upsilon(4S)$ in the Earth frame. We convert the time recording of each event occurrence to calculate GMST (Greenwich Mean Sidereal Time) as specified by the U.S. Naval Observatory. Figure 2 shows the A_{CPT} asymmetry, integrated over the Δt variable, as a function of the GMST.




FIG. 2: The opposite-sign lepton asymmetry A_{CPT} as a function of GMST (in seconds) folded over a period of 24 sidereal hours. The curve is a projection from the two-dimensional likelihood fit onto the sidereal time axis.

The results for sidereal-time dependent CPT violation

parameters are

$$\begin{aligned}\text{Im } z_0 &= (-14.1 \pm 7.3_{(\text{stat.})} \pm 2.4_{(\text{syst.})}) \times 10^{-3}, \\ \Delta\Gamma \cdot \text{Re } z_0 &= (-7.2 \pm 4.1_{(\text{stat.})} \pm 2.1_{(\text{syst.})}) \times 10^{-3} \text{ ps}^{-1}, \\ \text{Im } z_1 &= (-24.0 \pm 10.7_{(\text{stat.})} \pm 5.9_{(\text{syst.})}) \times 10^{-3}, \\ \Delta\Gamma \cdot \text{Re } z_1 &= (-18.8 \pm 5.5_{(\text{stat.})} \pm 4.0_{(\text{syst.})}) \times 10^{-3} \text{ ps}^{-1}.\end{aligned}$$

III. THE PARTIAL RECONSTRUCTION METHOD

In a complementary approach [8] the partial reconstruction of the $B^0 \rightarrow D^{*-}\ell^+\nu_\ell$ decay of one of the neutral B mesons is exploited. Though the available statistics is not as high as in the dilepton case, we can keep the charged B background at a lower level. At the same time, since the reconstructed and tag side are well defined, a procedure to determine particle detection asymmetries from data can be carried out. In this way, no dedicated control samples are needed to determine the asymmetry induced by the experimental cuts. A *BABAR* data sample equivalent to an integrated luminosity of 201 fb^{-1} at the $\Upsilon(4S)$ resonance (220 million $B\bar{B}$ pairs) is used for this study.

Preliminary cuts for lepton selection and non- $B\bar{B}$ background rejection are similar to the inclusive dilepton analysis. Only the charged lepton from the B^0 decay and the soft pion (π_s^-) from the D^{*-} decay are reconstructed. The $\bar{D}^0$ decay is not reconstructed, resulting in high selection efficiency. Due to the limited phase space available in the D^* decay, we approximate the direction of the D^* to be that of the π_s and estimate the energy $\tilde{E}_{D^*}$ of the D^* as a linear function of the energy of the π_s . We define the square of the missing neutrino mass as:

$$M_\nu^2 = \left(\sqrt{s}/2 - \tilde{E}_{D^*} - E_\ell \right)^2 - (\tilde{\mathbf{p}}_{D^*} + \mathbf{p}_\ell)^2, \quad (7)$$

where all quantities are defined in the $\Upsilon(4S)$ frame. We neglect the momentum of the B^0 (approximately $0.34 \text{ GeV}/c$), and identify the B^0 energy with half the total energy of the events ($\sqrt{s}/2$). E_ℓ and $\mathbf{p}_\ell$ are the energy and momentum vector of the lepton and $\tilde{\mathbf{p}}_{D^*}$ is the estimated momentum vector of the D^* . The distribution of M_ν^2 is peaked for signal events, while it is spread over a wide range for background events. Backgrounds are also suppressed using cuts on the lepton momenta, on the soft pion momentum, on the vertex fit probability, on $|\Delta t|$ and $\sigma_{\Delta t}$.

These selection criteria accept 470,877 events in the data sample (*tagged* events), and 5,291,868 partially reconstructed events that fail only the requirements of lepton tagging (*untagged* events). Fig. 3 shows the distributions of the squared neutrino mass for tagged events. Different components are recognized in the total sample: *signal* events, due to $D^{*-}\ell^+\nu_\ell$ events; *peaking* background events, that correspond to decays of charged B

mesons which peak in the M_ν^2 distribution, like $B^+ \rightarrow D^{*0}\ell^+\nu_\ell$, $D^{*0} \rightarrow D^{*-}\pi^+$; *combinatorial* events, corresponding to all $B\bar{B}$ decays not included in the previous categories; *continuum* events, coming from lepton-pair and light quark decays. Fractions of the different components in data are determined by means of fits to the M_ν^2 distribution: we obtain a signal fraction of $(46.6 \pm 0.9)\%$ in the selected mass region $-4.0 < M_\nu^2 < 2.0 \text{ GeV}^2/c^4$.




FIG. 3: Squared neutrino mass distributions for the tagged data events. The following fitted contributions are superimposed: continuum (solid grey line), combinatorial (dotted), $B^\pm$ peaking (dash-dotted), signal (dashed), all (solid black).

As in the dilepton case, the contributions to our sample are modeled using probability density functions (PDF) in Δt , based on the signal-background categories defined above and on the true origin of the tagging lepton.

An original method is used to determine possible charge detection asymmetries in the sample. We parameterize the total number of expected events for each tagged and untagged category in a likelihood function, which contains a term coming from the shapes of the PDFs for tagged events, an extended term for the number of tagged events and an extended term for the number of untagged events. This method reduces the correlation between CP mixing asymmetry and particle detection asymmetry.

Figure 4 shows the result of the maximum likelihood fit in terms of the projected asymmetries $A_{CP/T}$ between mixed positive and negative events, that we convert in a value for $|q/p|$. The main systematic uncertainties comes from the $B\bar{B}$ background and continuum charge asymmetries and the signal fraction determination from the M_ν^2 fit. The preliminary result is:

$$|q/p| - 1 = (6.5 \pm 3.4_{(\text{stat.})} \pm 2.0_{(\text{syst.})}) \times 10^{-3}, \quad (8)$$

which is compatible with the absence of CP violation in mixing at the 9.8% C.L.

IV. CONCLUSIONS

We have presented determinations of the parameters $|q/p|$, that is related to CP violation in $B^0\bar{B}^0$ mixing, and $\text{Im } z$, $\Delta\Gamma \cdot \text{Re } z$, $\text{Im } z_1$ and $\Delta\Gamma \cdot \text{Re } z_1$, that measure




FIG. 4: Raw asymmetry between mixed positive ($s_r = 1, s_t = 1$) and mixed negative ($s_r = -1, s_t = -1$) events with the PDF asymmetry derived from the fit superimposed. The asymmetry at low values of $|\Delta t|$ corresponds to the kaon-mistag contribution in decay-side tagged events.

CPT violation in a theory that takes into account effects of the implied Lorentz violation.

We exploit fits to Δt , the time difference between the two B decays in two complementary approaches. In the former, two high-momentum leptons are searched for in order to select inclusive semileptonic B^0 decays. In the latter, one of the B mesons is partially reconstructed in the semileptonic channel $D^{*-}\ell^+\nu_\ell$, i.e. only the lepton and the soft pion from $D^{*-} \rightarrow \bar{D}^0\pi^-$ decay are reconstructed, while the flavor of the other B is determined by means of lepton tagging. We use $B^0\bar{B}^0$ pair events, collected by the *BABAR* detector in the period 1999-2004.

No evidence is found of CP violation in mixing with either of the two methods: both results are compatible with the SM expectations and with previously published *BABAR* results [9, 10]. The experimental values of $A_{CP/T}$ are used as constraints [11], together with other measurements related to the unitarity triangle, in a global fit to the CKM theory parameters that also includes the possibility of NP. Figure 5 shows the most recent results of this fit in terms of the allowed regions for the parameters C_d and ϕ_d defined in Eq. (4), corresponding to a 95% confidence interval of $[0.59, 2.51]$ for C_d and $[-7.0, 1.0]^\circ$

for ϕ_d .




FIG. 5: Allowed regions for the NP parameters C_d and ϕ_d at the 68% (dark blue) and 95% (light blue) confidence levels.

For the $\text{Im}z$, $\Delta\Gamma \cdot \text{Re}z$ pair, the results are compatible with the SM expectation (0,0) within 1.5σ . When sidereal-time dependence is taken into account, the result in the $\text{Im}z_1$ - $\Delta\Gamma \cdot \text{Re}z_1$ plane has a significance of 2.2σ , compatible with the absence of CPT violation in mixing (Figure 6).




FIG. 6: Allowed regions for the CPT -violating parameters $\text{Im}z_1$ and $\Delta\Gamma \cdot \text{Re}z_1$ at various confidence levels. The red star represents the SM expectation and the solid black ellipses correspond to 1, 2 and 3σ significances.

[1] M. Ciuchini *et al.*, JHEP **0308**, 031 (2002).
[2] M. Beneke *et al.*, Phys. Lett. **B 576**, 173 (2003).
[3] S. Laplace *et al.*, Phys. Rev. D **65**, 094040 (2002).
[4] M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
[5] V. A. Kostelecký, Phys. Rev. Lett. **80**, 1818 (2004).
[6] B. Aubert *et al.*, The *BABAR* Collaboration, Phys. Rev. Lett. **96**, 251802 (2006).
[7] B. Aubert *et al.*, The *BABAR* Collaboration, NIM A **479** (2002) 1.

[8] B. Aubert *et al.*, The *BABAR* Collaboration, contributed to 33rd International Conference on High Energy Physics (ICHEP 06), Moscow, Russia [hep-ex/0607091].
[9] B. Aubert *et al.*, The *BABAR* Collaboration, Phys. Rev. **D70**, 012007 (2004).
[10] B. Aubert *et al.*, The *BABAR* Collaboration, Phys. Rev. Lett. **88**, 231801 (2002).
[11] M. Bona *et al.*, The *UTfit* Collaboration, hep-ph/0605213.